

Large Acute Spinal Posterior Subdural Hematoma Complicating Thrombolysis with Streptokinase in Acute Myocardial Infarction

Wahane M¹, Chopade S², Janbandhu A³

ABSTRACT

Thrombolysis therapy is the cornerstone of treatment of ST elevation myocardial infarction. It is associated with multiple complications. One of the complications is intracranial bleeding. But intraspinal hemorrhage causing cord compression is rare. Still rarer is subdural posterior spinal hematoma resulting in quadriplegia. We here present a case of subdural posterior spinal hematoma following Thrombolytic therapy in Acute MI

Key words : Spinal subdural hematoma, Streptokinase, Thrombolysis

Introduction :

Thrombolysis therapy is the cornerstone of treatment of ST elevation myocardial infarction. It is associated with multiple complications. One of the complications is intracranial bleeding. But intraspinal hemorrhage causing cord compression is rare. Still rarer is subdural posterior spinal hematoma resulting in quadriplegia. We here report a case of subdural posterior spinal hematoma resulting in quadriplegia following thrombolytic therapy in acute myocardial infarction.

Case Report :

35 year old male patient came to the hospital with chief complaints of Retrosternal chest pain & sweating of 2 hours duration prior to hospitalization without any other significant presentsymptoms. He was non diabetic, non hypertensive without any relevant past history.

His physical examination revealed pulse of 56/min regular, BP 90/60 mmhg & no evidence of congestive heart failure Systemic examination did not reveal any abnormality. On investigation ECG showed -2 mm ST elevation in II, III and aVF leads, -ST depression with T wave inversion in I and aVL,

V2, - Right sided lead showed ST elevation in V4R. Suggestive of Inferior wall myocardial infarction with RV infarct. (**Fig. 1**) CPKMB was raised.

The patient was given fluid challenge with 200 mL Normal Saline and thrombolysed with 1.5 MU STK. Following the thrombolysis, the chest pain decreased, heart rate increased to 78/min, regular.

Blood pressure increased to 120/80 mmHg. ECG after thrombolysis showed decrease in ST elevation.

Patient tolerated thrombolysis well without any arrhythmias or other immediate complications.

Routine investigations were normal as follows :

Sr. No.	Investigation	Result
1	Hemoglobin	10 g%,
2	Total leukocyte count	10,000/mm ³
3	Serum creatinine	1.2 mg/dl
4	Blood urea level	27 mg/dl
5	INR	1.3

Apart from Thrombolysis he was treated with antiplatelets, statins, ACE inhibitor & beta blocker after BP improved. Patient was stable for 36 hrs.

36 hours after thrombolysis the patient complained of sudden onset severe back pain in upper thoracic region which increased on coughing & developed weakness in both lower limbs and retention of urine. In the next 4 hours, patient developed profound weakness in all 4 limbs. Neurological examination revealed acute onset, progressive sensory motor quadriplegia with bladder involvement.

¹Assistant Professor, ²Resident
Department of Medicine, IGGMC, Nagpur
³Consultant Physician, Yavatmal

Address for Correspondence -

Dr. Madhuri Wahane

E-mail :

MRI cervical and thoracic spine revealed, large acute extra axial intraspinal posterior subdural hematoma from C2 to D3 level causing cord compression and intramedullary hyperintense signals seen in cord from C7 to D3 levels suggestive of cord edema. (*Fig.2 & 3*)

Antiplatelet drugs were stopped. Coagulation profile was immediately obtained. Patient was referred to neurosurgeon for decompression. Patient was operated immediately and evacuation of hematoma was done. The patient gradually improved and had complete recovery in the next 7 days.

During follow up, patient had no angina, no features of heart failure and no neurological deficit.

Discussion :

Streptokinase therapy for acute myocardial infarction is associated with various bleeding complications. Intracranial hemorrhage, occurs in 0.46-0.88% of patients treated with thrombolytic agents.¹ Unlike intracranial hemorrhage, intraspinal hemorrhage usually occurs in epidural space most often in dorsal thoracic spine. Incidence of spinal subdural hematoma after thrombolysis is extremely rare.²

The most frequent clinical symptom of a intraspinal hematoma is neck or back pain with or without radicular symptoms. Neurological deficits from spinal cord compression can appear insidiously or abruptly and progress in a wide clinical picture of complete paraplegia/quadriplegia to Brown Séquard's syndrome.³

MRI scan is the investigation of choice for suspected intraspinal hematoma. It provides multiplanar accurate information on both the location and extent of hematoma, as well as the intensity of spinal cord compression by the lesion and is useful in differentiating accompanying intraspinal masses and to follow-up the resolution of the hematoma.⁴

Emergency surgery is the treatment preferred for intraspinal hemorrhages but patient-specific factors should be carefully evaluated prior to surgery. Anticoagulants and thrombolytics should be

immediately discontinued and correction of coagulopathy should be undertaken prior to surgery.⁵



Fig. 2 : Sagittal section of spinal cord showing posterior subdural spinal hematoma




Fig. 3 : Transverse section of spinal cord showing posterior subdural spinal hematoma

Conservative treatment is still an important option of treatment in some selective patients with mild and rapidly spontaneous recovery symptoms or high surgical risk patients with bleeding tendency associated with severe systemic disease, advanced cardiovascular disease or advanced and irreversible spinal cord injury. Patients can be managed conservatively with reversal of coagulant effects, close observation of neurologic deficits and in occasional cases, methylprednisolone administration may achieve good results without surgery.⁶

Conclusion :

Thrombolytic therapy has significant role in reperfusion for patients of acute myocardial infarction

Presenting within 12 hours. The physician should be careful in monitoring the patients, as it may result in bleeding complications. Intraspinial hemorrhage is rare complication, early recognition & management is crucial in preventing neurodeficit.

Conflicts of interest : none reported by authors

References :

1. Ozgosmen S, Yoldas T, Kocakoc E *et al.* Spinal epidural hematoma associated with streptokinase treatment for myocardial infarction. *Spinal cord* 2004; 42: 374-377.
2. Dahlin PA, George J : Intraspinial hematoma as a complication of anticoagulant therapy; *Clin Pharm.* 1984 Nov-Dec ;3(6):656-61.
3. Kreppel D, Antoniadis G, Seeling W. Spinal hematoma : a literature survey with a meta-analysis of 613 patients. *Neurosurg Rev* 2003; 26: 149.
4. vanHeesewijk JPM, Caparie JWBM. Acute spontaneous spinal epidural hematoma in a child. *EurRadiol* 2000; 10:1874-1876.
5. Connolly SE, Winfree CJ, McCormick PC. Management of spinal epidural hematoma after tissue plasminogen activator. A case report. *Spine* 1996; 21: 1694-1698.
6. Hentschel SJ, Woolfenden AR, Fairholm DJ. Resolution of spontaneous spinal epidural hematoma without surgery. *Spine* 2001; 26:E525-E527.